

REMARKS

The present Preliminary Amendment is submitted to delete the multiple dependency of the claims, thereby placing such claims in condition for examination and reducing the required PTO filing fee.

Attached hereto is a marked-up version of the changes made to the specification and claims by the current Amendment. The attached page is captioned "Version With Markings to Show Changes Made".

Respectfully submitted,

Yoshiro SHIOKAWA

By



Nils E. Pedersen  
Registration No. 33,145  
Attorney for Applicant

NEP/krl  
Washington, D.C. 20006-1021  
Telephone (202) 721-8200  
Facsimile (202) 721-8250  
April 3, 2001

THE COMMISSIONER IS AUTHORIZED  
TO CHARGE ANY DEFICIENCY IN THE  
FEES FOR THIS PAPER TO DEPOSIT  
ACCOUNT NO. 23-0975

Version with Markings to  
Show Changes Made

6. (Amended) The Q-pole type mass spectrometer according to [any one of claims 2 to 5] claim 2, characterized in that upon control on the motion of ion to be measured in the axial direction within the Q-pole region, Coulomb force is employed, the said Coulomb force is generated by electric field formed by four Q-poles composing the Q-pole type mass spectrometer, so constructed that four Q-poles have an equal DC potentials except DC voltage: U at the same position in the axial direction of each Q-pole of four Q-poles, while the each Q-pole of four Q-poles has different DC potentials depending on their positions in the axial direction.

8. (Amended) The Q-pole type mass spectrometer according to [any one of claims 2 to 5] claim 2, characterized in that control on the motion of ion to be measured in the axial direction within the Q-pole region uses a reaction force generated by a collision between the ion to be measured and the atmospheric gas.

10. (Amended) The Q-pole type mass spectrometer according to [any one of claims 2 to 5] claim 2, characterized in that control on the motion of ion to be measured in the axial direction within the Q-pole region is carried out by setting the length of the Q-pole, kind and pressure of the atmospheric gas, potential of the ion source and potential on the axis of the Q-pole so that the ion to be measured is capable of passing the Q-pole region without receiving any additional force in the axial direction.

11. (Amended) The Q-pole type mass spectrometer according to [any one of claims 2 to 5] claim 2, characterized in that control on the motion of ion to be measured in the axial direction within the Q-pole region is carried out using Coulomb force generated by space charge formed by the ion to be measured within the Q-pole region.

13. (Amended) The Q-pole type mass spectrometer according to [any one of claims 2 to 5] claim 2, characterized in that control on the motion of ion to be measured in the axial direction within the Q-pole region is carried out using Lorentz force generated by high-frequency magnetic field synchronous with quadrupole high-frequency electric field applied in the diameter direction.

14. (Amended) The Q-pole type mass spectrometer according to [any one of claims 2 to 5] claim 2, characterized in that control on the motion of ion to be measured in the axial direction within the Q-pole region is carried out using electromagnetic induction force generated by a magnetic field changing in its intensity with time passage, applied in the diameter direction.